

On the Obstruction to the Exactness of the Variational Bicomplex

G.GIACHETTA^{†1}, L.MANGIAROTTI^{†2} AND G.SARDANASHVILY^{‡3}

[†] *Department of Mathematics and Physics, University of Camerino, 62032 Camerino (MC), Italy*

[‡] *Department of Theoretical Physics, Physics Faculty, Moscow State University, 117234 Moscow, Russia*

Abstract. We study the obstruction to the exactness of the variational complex for a field theory on an affine bundle.

Mathematics Subject Classification (2000). 53C80, 57R22, 58E35, 70S05.

1 Introduction

Let $Y \rightarrow X$ be a real smooth fibre bundle coordinated by (x^λ, y^i) , and Ω_∞^* the algebra of exterior forms on all finite order jet manifolds of $Y \rightarrow X$ modulo the pull-back identification. Put $\dim X = n$. We have the variational complex

$$0 \rightarrow \mathbf{R} \rightarrow \Omega_\infty^0 \xrightarrow{d_H} \Omega_\infty^{0,1} \xrightarrow{d_H} \dots \xrightarrow{d_H} \Omega_\infty^{0,n} \xrightarrow{\varepsilon_1} E_1 \xrightarrow{\varepsilon_2} E_2 \longrightarrow \dots, \quad (1)$$

where $\Omega_\infty^{0,*}$ is a subalgebra of horizontal (semibasic) exterior forms, d_H is the horizontal exterior differential, ε_1 is the Euler–Lagrange map, ε_2 is the Helmholtz–Sonin map, and so on [3, 4, 6, 7, 8, 14, 15]. This complex provides the algebraic approach to the calculus of variations in field theory. The well-known theorem states the local exactness of this complex as follows.

THEOREM 1. *If a fibre bundle Y is $\mathbf{R}^{n+m} \rightarrow \mathbf{R}^n$, the variational complex (1) except the first term is exact, i.e.*

$$(i) \text{ Ker } d_H = \text{Im } d_H, \quad (ii) \text{ Ker } \varepsilon_1 = \text{Im } d_H, \quad (iii) \text{ Ker } \varepsilon_{k+1} = \text{Im } \varepsilon_k.$$

REMARK 1. As in the De Rham complex on a connected manifold, one puts $H^0(d_H) = \mathbf{R}$ because, if $d_H f = 0$, $f \in \Omega_\infty^0$, then $f = \text{const}$. From now on, by H^* we will mean the cohomology groups of order $k > 0$.

¹giachetta@campus.unicam.it

²mangiaro@camserv.unicam.it

³sard@campus.unicam.it; sard@grav.phys.msu.su

Restricted to $\Omega_\infty^{0,*}$, the variational complex (1) yields the so called horizontal complex

$$0 \rightarrow \mathbf{R} \rightarrow \Omega_\infty^0 \xrightarrow{d_H} \Omega_\infty^{0,1} \xrightarrow{d_H} \dots \xrightarrow{d_H} \Omega_\infty^{0,n} \xrightarrow{d_H} 0. \quad (2)$$

Then a corollary of Theorem 1 is the algebraic Poincaré lemma on the local exactness of the complex (2).

PROPOSITION 2. *If a fibre bundle Y is $\mathbf{R}^{n+m} \rightarrow \mathbf{R}^n$, the horizontal complex has the cohomology groups*

$$H^{k < n}(d_H) = 0, \quad H^n(d_H) = \varepsilon_1(\Omega_\infty^{0,n}). \quad (3)$$

This work is devoted to the obstruction to the (global) exactness of the complexes (1) and (2).

It should be emphasized that we consider the variational bicomplex in the class of smooth exterior forms Ω_∞^* on the infinite-order jet space $J^\infty Y$ which are the pull-back of exterior forms on finite-order jets. This is not the case of [2, 3]. In these works smooth forms on $J^\infty Y$ which are locally the pull-back of forms on finite-order jet manifolds are considered. The key point is that $J^\infty Y$ admits a partition of unity with respect to this class of smooth functions. Then, applying the well-known Mayer-Vietoris sequence [5], one can show that contact d_H -closed forms are necessarily d_H -exact. This may not be true for the class of exterior forms Ω_∞^* which usually are utilized in field theory.

We show that, if a fibre bundle $Y \rightarrow X$ admits a global section (e.g., if it is an affine bundle), there is a monomorphism of the De Rham homology groups $H^*(X)$ of the base manifold X to the cohomology groups H_{var}^* of the variational complex (1) and to those $H^*(d_H)$ of the horizontal complex (2). When $Y \rightarrow X$ is a vector bundle, we find the obstruction to the exactness of the variational complex (1) at E_k . We show that, in this case, the cohomology group of the variational complex (1) at $\Omega_\infty^{0,n}$ is equal to the cohomology group $H^n(X)$ of the base manifold. It states the following theorem (see also [10]).

THEOREM 3. *If $Y \rightarrow X$ is an affine bundle, an r -order Lagrangian $L \in \Omega_r^{0,n}$ is variationally trivial iff it takes the form $L = d_H \sigma$ where $\sigma \in \Omega_{r-1}^{0,n-1}$ is a horizontal $(n-1)$ -form on the jet manifold $J^{r-1}Y$.*

Recall that the familiar result states that L is locally d_H -exact.

2 Preliminaries

Unless otherwise stated, manifolds throughout are assumed to be real, smooth, finite-dimensional, Hausdorff, paracompact and connected.

Let $J^r Y$ be the r -order jet manifold of sections of a fibre bundle $Y \rightarrow X$. It is endowed with the adapted coordinates (x^λ, y_Λ^i) , $0 \leq |\Lambda| \leq r$, where $\Lambda = (\lambda_k \dots \lambda_1)$, $|\Lambda| = k$, is a symmetric multi-index. By $\Lambda + \Sigma$ we will denote the symmetric multi-index $(\lambda_k \dots \lambda_1 \sigma_m \dots \sigma_1)$. The transition functions of these coordinates read

$$y_{\lambda+\Lambda}^i = \frac{\partial x^\mu}{\partial x'^\lambda} d_\mu y_\Lambda^i, \quad (4)$$

where by d_λ are meant the total derivatives

$$d_\lambda = \partial_\lambda + \sum_{|\Lambda|=0} y_{\Lambda+\lambda}^i \partial_i^\Lambda.$$

We will use the notation

$$\partial_\Lambda = \partial_{\lambda_k} \circ \dots \circ \partial_{\lambda_1}, \quad d_\Lambda = d_{\lambda_k} \circ \dots \circ d_{\lambda_1}, \quad \Lambda = (\lambda_k \dots \lambda_1).$$

There is the inverse system

$$X \xleftarrow{\pi} Y \xleftarrow{\pi_0^1} \dots \xleftarrow{\pi_{r-1}^r} J^r Y \xleftarrow{\pi_r^r} \dots \quad (5)$$

of fibrations (surjective submersions) of finite order jet manifolds. This inverse system has a projective limit $J^\infty Y$. It is a paracompact Fréchet manifold which admits a smooth partition of unity [1, 4, 13, 14]. Differential objects on $J^\infty Y$ are introduced as operations on the $\mathbf{R}$ -ring $C^\infty(J^\infty Y)$ of smooth locally cylindrical functions. A real function f on $J^\infty Y$ is said to be a smooth (locally cylindrical) function if it is locally a pull-back of a smooth function on some finite-order jet manifold $J^k Y$ with respect to the surjection $\pi_k^\infty : J^\infty Y \rightarrow J^k Y$. Vector fields on $J^\infty Y$ are defined as derivations of this ring. They make up the left locally free $C^\infty(J^\infty Y)$ -module $\text{Der}(C^\infty(J^\infty Y))$. The $C^\infty(J^\infty Y)$ -dual of $\text{Der}(C^\infty(J^\infty Y))$ is the $C^\infty(J^\infty Y)$ -module of exterior 1-forms on $J^\infty Y$. In field theory, one usually considers the direct limit Ω_∞^* in the category of $\mathbf{R}$ -modules of the direct system

$$\Omega^*(X) \xrightarrow{\pi^*} \Omega^*(Y) \xrightarrow{\pi_0^{1*}} \Omega_1^* \xrightarrow{\pi_1^{2*}} \dots \xrightarrow{\pi_{r-1}^{r*}} \Omega_r^* \longrightarrow \dots \quad (6)$$

of the $\mathbf{R}$ -modules Ω_k^* of exterior forms on finite-order jet manifolds $J^k Y$. It consists of pull-backs of exterior forms on finite-order jet manifolds.

The $\mathbf{R}$ -module Ω_∞^* possesses the structure of a filtered module given by the Ω_k^0 -submodules Ω_k^* of Ω_∞^* [9]. An endomorphism γ of Ω_∞^* is called a filtered morphism if there exists $i \in \mathbf{N}$ such that the restriction of γ to Ω_k^* is the homomorphism of Ω_k^* into Ω_{k+i}^* over the injection $\Omega_k^0 \hookrightarrow \Omega_{k+i}^0$ for all k .

THEOREM 4. [12]. *Every direct system of endomorphisms $\{\gamma_k\}$ of Ω_k^* such that*

$$\pi_k^{j*} \circ \gamma_k = \gamma_j \circ \pi_k^{j*}$$

for all $j > k$ has the direct limit γ_∞ in filtered endomorphisms of Ω_∞^* . If all γ_k are monomorphisms [epimorphisms], then γ_∞ is also a monomorphism [epimorphism].

By virtue of this theorem, the standard algebraic operations on exterior forms on finite-order jet manifolds have the direct limits on Ω_∞^* , and make Ω_∞^* an exterior algebra.

Since Ω_∞^* consists of pull-back forms, its elements can be written in the familiar coordinate form. In particular, the basic 1-forms dx^λ and the contact 1-forms $\theta_\Lambda^i = dy_\Lambda^i - y_{\Lambda+\Lambda}^i dx^\lambda$ provide the local generators of Ω_∞^* , and define the canonical splitting of the space of m -forms

$$\Omega_\infty^m = \Omega_\infty^{0,m} \oplus \Omega_\infty^{1,m-1} \oplus \dots \oplus \Omega_\infty^{m,0}, \quad (7)$$

in the spaces $\Omega_\infty^{k,m-k}$ of k -contact forms. We have the corresponding k -contact projections

$$h_k : \Omega_\infty^m \rightarrow \Omega_\infty^{k,m-k}, \quad 1 \leq k \leq m,$$

and the horizontal projection

$$h_0 : \Omega_\infty^m \rightarrow \Omega_\infty^{0,m}.$$

Accordingly, the exterior differential on Ω_∞^* is decomposed into the sum $d = d_H + d_V$ of horizontal and vertical differentials

$$\begin{aligned} d_H : \Omega_\infty^{k,s} &\rightarrow \Omega_\infty^{k,s+1}, & d_H(\phi) &= dx^\lambda \wedge d_\lambda(\phi), & \phi &\in \Omega_\infty^*, \\ d_V : \Omega_\infty^{k,s} &\rightarrow \Omega_\infty^{k+1,s}, & d_V(\phi) &= \theta_\Lambda^i \wedge \partial_\Lambda^i \phi. \end{aligned}$$

They obey the nilpotency rule

$$d_H \circ d_H = 0, \quad d_V \circ d_V = 0, \quad d_V \circ d_H + d_H \circ d_V = 0 \quad (8)$$

and the relations

$$h_0 d = d_H h_0, \quad (9)$$

$$h_k d h_k = d_H h_k, \quad (10)$$

3 Cohomology of the infinite-order De Rham complex

The exterior algebra Ω_∞^* provides the infinite-order De Rham complex

$$0 \longrightarrow \mathbf{R} \longrightarrow \Omega_\infty^0 \xrightarrow{d} \Omega_\infty^1 \xrightarrow{d} \dots \quad (11)$$

PROPOSITION 5. *The cohomology $H^*(\Omega_\infty^*)$ of the infinite-order De Rham complex (11) coincides with the De Rham cohomology $H^*(Y)$ of the fibre bundle Y .*

The result is a corollary of the following two assertions.

PROPOSITION 6. [12]. *The operation of taking cohomology groups of a cochain complex commutes with the passage to the direct limit.*

It is a corollary of Theorem 4.

PROPOSITION 7. [4]. *The De Rham cohomology $H^*(J^k Y)$ of any finite-order jet manifold $J^k Y$ is equal to that of the fibre bundle Y .*

It is based on the fact that jet bundles $J^k Y \rightarrow J^{k-1} Y$ are affine, while the De Rham cohomology of an affine bundle is equal to that of its base.

Recall that, given a fibre bundle $\pi : Y \rightarrow X$, there is a homomorphism

$$\pi^* : H^*(X) \rightarrow H^*(Y) \quad (12)$$

of the De Rham cohomology groups of its base X to those of Y .

LEMMA 8. *If a fibre bundle $Y \rightarrow X$ admits a global section s , there is the sequence of homomorphisms of cohomology groups*

$$H^*(X) \xrightarrow{\pi^*} H^*(Y) \xrightarrow{s^*} H^*(X), \quad (13)$$

where s^* is an epimorphism and, consequently, π^* (12) is a monomorphism.

If $Y \rightarrow X$ admits a global section, we have the corresponding monomorphism

$$\pi^* : H^*(X) \hookrightarrow H^*(\Omega_\infty^*) \quad (14)$$

of the De Rham cohomology groups of the base X to the cohomology groups of the infinite-order De Rham complex (11).

If $Y \rightarrow X$ is an affine bundle, the monomorphism (14) is an isomorphism

$$H^*(\Omega_\infty^*) = H^*(X). \quad (15)$$

In this case, any closed form $\phi \in \Omega_\infty^*$ is decomposed into the sum

$$\phi = \varphi + d\xi \quad (16)$$

where $\varphi \in \Omega^*(X)$ is a closed form on X .

4 Cohomology of the horizontal complex

Due to the nilpotency rule (8), the vertical and horizontal differentials d_V and d_H define the bicomplex

$$\begin{array}{ccccccc}
 & & \uparrow & & \uparrow & & \\
 \dots & \longrightarrow & \Omega_{\infty}^{k+1,m} & \xrightarrow{d_H} & \Omega_{\infty}^{k+1,m+1} & \longrightarrow & \dots \\
 & & \uparrow d_V & & \uparrow d_V & & \\
 \dots & \longrightarrow & \Omega_{\infty}^{k,m} & \xrightarrow{d_H} & \Omega_{\infty}^{k,m+1} & \longrightarrow & \dots \\
 & & \uparrow & & \uparrow & &
 \end{array} \tag{17}$$

The rows and columns of these bicomplex are horizontal and vertical complexes. Let us start from the vertical ones.

PROPOSITION 9. *If $Y \rightarrow X$ is a vector bundle, then any d_V -closed form ϕ is the sum $\phi = d_V\sigma + \varphi$ of a d_V -exact form and an exterior form on X , i.e., vertical complexes are exact.*

Proof. Local exactness of a vertical complex on a coordinate chart (x^λ, y^i) follows from a version of the Poincaré lemma with parameters. We have the the corresponding homotopy operator

$$\sigma = \int_0^1 t^k [\bar{y}] \phi(x^\lambda, ty_\lambda^i) dt, \quad \phi \in \Omega^{k,*}, \tag{18}$$

where $\bar{y} = y_\lambda^i \partial_i^\lambda$. Since $Y \rightarrow X$ is a vector bundle, it is readily observed that the homotopy operator is globally defined, and so is the form σ . $\square$

Turn now to the rows of the diagram (17). We will start from the horizontal complex (2). Due to the relation (9), the horizontal projection h_0 defines a homomorphism of the infinite-order De Rham complex (11) to the horizontal complex (2) which sends closed and exact forms to d_H -closed and d_H -exact horizontal forms, respectively. Accordingly, it yields the homomorphism of the cohomology groups of these complexes

$$h_0^* : H^*(\Omega_{\infty}^*) \rightarrow H^*(d_H), \tag{19}$$

which is neither monomorphism nor epimorphism in general. By virtue of Proposition 5, the homomorphism (19) is the homomorphism of the cohomology groups

$$h_0^* : H^*(Y) \rightarrow H^*(d_H). \tag{20}$$

Then we also have the homomorphism

$$h_0^* \circ \pi^* : H^*(X) \rightarrow H^*(d_H). \tag{21}$$

PROPOSITION 10. *If a fibre bundle $Y \rightarrow X$ admits a global section s , the homomorphism (21) is a monomorphism.*

Proof. Bearing in mind the monomorphism (14), let $\phi \in \Omega_\infty^*$ be a closed form whose cohomology class belongs to $H^*(X) \subset H^*(\Omega_\infty^*)$. We will show that, if $h_0\phi$ is d_H -exact, ϕ is exact. The form ϕ is represented by the sum (16), and we have

$$h_0\phi = \varphi + d_H(h_0\xi).$$

Let $\varphi = d_H\sigma$ where σ is an exterior form on some finite-order jet manifold J^kY . Given a global section s of the fibre bundle $Y \rightarrow X$, let $J^{k+1}s$ be its $(k+1)$ -order jet prolongation to a section of the jet bundle $J^{k+1}Y \rightarrow X$. Then it is readily observed that

$$\varphi = (J^{k+1}s)^*(d_H\sigma) = (J^{k+1}s)^*(d\sigma) = d((J^{k+1}s)^*\sigma).$$

□

Since h_0 is a homomorphism of the infinite-order De Rham complex (11) to the horizontal complex (2), we have the simple exact sequence of the cochain complexes

$$\begin{array}{ccccccc} & 0 & & 0 & & 0 & \\ & \downarrow & & \downarrow & & \downarrow & \\ \cdots & \xrightarrow{d} & \text{Ker } h_0 & \xrightarrow{d} & \text{Ker } h_0 & \xrightarrow{d} \cdots & \xrightarrow{d} \text{Ker } h_0 \xrightarrow{d} \text{Ker } h_0 \\ & \text{in} \downarrow & & \text{in} \downarrow & & \text{in} \downarrow & \parallel \\ \cdots & \xrightarrow{d} & \Omega_\infty^{k-1} & \xrightarrow{d} & \Omega_\infty^k & \xrightarrow{d} \cdots & \xrightarrow{d} \Omega_\infty^n \xrightarrow{d} \Omega_\infty^{n+1} \longrightarrow \cdots \\ & h_0 \downarrow & & h_0 \downarrow & & h_0 \downarrow & \downarrow \\ \cdots & \xrightarrow{d_H} & \Omega_\infty^{0,k-1} & \xrightarrow{d_H} & \Omega_\infty^{0,k} & \xrightarrow{d_H} \cdots & \xrightarrow{d_H} \Omega_\infty^{0,n} \longrightarrow 0 \\ & \downarrow & & \downarrow & & \downarrow & \downarrow \\ & 0 & & 0 & & 0 & \end{array} \quad (22)$$

Its first row is a subcomplex of the infinite-order De Rham complex (11). There is the corresponding exact sequence of cohomology groups

$$\cdots \longrightarrow H^k(\text{Ker } h_0) \xrightarrow{\text{in}^*} H^k(\Omega_\infty^*) \xrightarrow{h_0^*} H^k(d_H) \xrightarrow{d^*} H^{k+1}(\text{Ker } h_0) \longrightarrow \cdots \quad (23)$$

We can say something more on this exact sequence when $Y \rightarrow X$ is an affine bundle.

PROPOSITION 11. *If $Y \rightarrow X$ is an affine bundle, the exact sequence of cohomology groups (23) takes the form*

$$\begin{aligned} \cdots \longrightarrow H^{k-1}(d_H)/H^{k-1}(X) &\xrightarrow{\text{in}^*} H^k(X) \xrightarrow{h_0^*} H^k(d_H) \xrightarrow{d^*} H^k(d_H)/H^k(X) \longrightarrow \\ \cdots \xrightarrow{\text{in}^*} H^n(X) &\xrightarrow{h_0^*} H^n(d_H) \xrightarrow{d^*} H^n(d_H)/H^n(X) \longrightarrow 0, \end{aligned} \quad (24)$$

where in^* is a homomorphism to $0 \in H^*(X)$ and h_0^* is a monomorphism.

Proof. Firstly, let us describe the cohomology group $H^k(\text{Ker } h_0)$ of the first row in the diagram (22). In this complex, the cocycles are closed forms $\phi \in \text{Ker } h_0$, while the coboundaries are exact forms $d\xi$ where $\xi \in \text{Ker } h_0$. Because of the isomorphism (15), such a k -cocycle is a coboundary in the infinite-order De Rham complex and is decomposed into the sum

$$\phi = d\sigma + d\xi, \quad (25)$$

where $\sigma \in \Omega_\infty^{0,k-1}$ is a horizontal form such that $h_0 d\sigma = 0$. It follows that $d_H \sigma = 0$, i.e., σ is a d_H -closed form. At the same time, if σ is a d_H -exact form $\sigma = d_H \psi$, then

$$d\sigma = dd_H \psi = -dd_V \psi$$

is a coboundary in $\text{Ker } h_0$. Therefore, we have an epimorphism of cohomology groups

$$d^* : H^{k-1}(d_H) \rightarrow H^k(\text{Ker } h_0).$$

Obviously, its kernel $\text{Ker } d^*$ contains the cohomology group $H^{k-1}(X) \subset H^{k-1}(d_H)$. We show that $\text{Ker } d^* = H^{k-1}(X)$. Let $\sigma \in \text{Ker } d^*$, i.e.,

$$d_H \sigma = 0, \quad d\sigma = \psi, \quad \psi \in \text{Ker } h_0.$$

Then we have $d(\sigma - \psi) = 0$, and

$$\sigma = \varphi + d\xi + \psi, \quad \varphi \in \Omega^*(X),$$

in accordance with the decomposition (16). Moreover, being horizontal,

$$\sigma = \varphi + h_0 d\xi = \varphi + d_H h_0 \xi,$$

i.e., it belongs to the d_H -cohomology class of the form φ on X . At last, since all cocycles in $\text{Ker } h_0$ are coboundaries in Ω_∞^* , the morphism in^* sends the cohomology group $H^k(\text{Ker } h_0)$ to zero. $\square$

Turn now to cohomology of the k -contact horizontal complex

$$\dots \xrightarrow{d_H} \Omega_\infty^{k,m-1} \xrightarrow{d_H} \Omega_\infty^{k,m} \xrightarrow{d_H} \dots \quad (26)$$

It is readily observed that, because of the relation (9), the contact projection h_k fails to be a homomorphism of the infinite-order De Rham complex (11) to the horizontal complex (26). Accordingly, we have no homomorphism of the corresponding cohomology groups. At the same time, due to the nilpotency rule (8), the horizontal differential yields

a homomorphism of the k -contact horizontal complex (26) to the $(k + 1)$ -contact one, together with the corresponding homomorphism of their cohomology groups

$$d_V^* : H^*(k, d_H) \rightarrow H^*(k + 1, d_H).$$

Moreover, we have the complex of the cohomology groups

$$0 \longrightarrow H^*(X) \xrightarrow{h_0^*} H^*(d_H) \xrightarrow{d_V^*} \cdots \xrightarrow{d_V^*} H^*(k, d_H) \xrightarrow{d_V^*} \cdots . \quad (27)$$

Its cohomology is called the second cohomology [11].

Similarly, the horizontal differential d_H defines homomorphisms of the vertical complexes.

Let us show that, if the fibre bundle $Y \rightarrow X$ admits a global section, the complex (27) is exact at the second term, i.e. the kernel of d_V in $H^*(d_H)$ coincides with the De Rham cohomology $H^*(X)$.

LEMMA 12. *If $\phi \in \text{Ker } d_H d_V$, then*

$$\phi = \sigma + \xi + \varphi, \quad (28)$$

where σ is a d_H -closed form, ξ is a d_V -closed form and $\varphi \in \Omega^*(X)$ (see [15] for the local case).

PROPOSITION 13. *If the fibre bundle $Y \rightarrow X$ admits a global section, the complex (27) is exact at the second term.*

Proof. Let $\phi \in \Omega_\infty^{0,*}$ be a d_H -closed form, i.e., $d_H \phi = 0$. It belongs to $\text{Ker } d_V^*$ if $d_V \phi = d_H \varepsilon$. In accordance with the relation (28), it follows that $\phi = d_H \varepsilon + \varphi$, i.e. belongs to $H^*(X) \subset H^*(d_H)$. $\square$

5 Cohomology of the variational complex

Obviously, the cohomology groups $H_{\text{var}}^{<n}$ of the variational complex (1) coincide with those of the horizontal complex (2). To say something on other cohomology groups $H_{\text{var}}^{\geq n}$, let us

consider the simple exact sequence of cochain complexes

$$\begin{array}{ccccccc}
& 0 & & 0 & & 0 & & 0 \\
& \downarrow & & \downarrow & & \downarrow & & \downarrow \\
\cdots \longrightarrow & \text{Ker } h_0 & \xrightarrow{d} & \text{Ker } h_0 & \xrightarrow{d} & \text{Ker } e_1 & \xrightarrow{d} & \text{Ker } e_2 \longrightarrow \cdots \\
& \downarrow & & \downarrow & & \downarrow & & \downarrow \\
\cdots \longrightarrow & \Omega_\infty^{n-1} & \xrightarrow{d} & \Omega_\infty^n & \xrightarrow{d} & \Omega_\infty^{n+1} & \xrightarrow{d} & \Omega_\infty^{n+2} \longrightarrow \cdots \\
& \downarrow h_0 & & \downarrow h_0 & & \downarrow e_1 & & \downarrow e_2 \\
\cdots \longrightarrow & \Omega_\infty^{0,n-1} & \xrightarrow{d_H} & \Omega_\infty^{0,n} & \xrightarrow{\varepsilon_1} & E_1 & \xrightarrow{\varepsilon_2} & E_2 \longrightarrow \cdots \\
& \downarrow & & \downarrow & & \downarrow & & \downarrow \\
& 0 & & 0 & & 0 & & 0
\end{array} \tag{29}$$

where $e_k = \tau_k \circ h_k$ and τ_k is the projection map providing the decomposition

$$\Omega_\infty^{k,n} = \tau_k(\Omega_\infty^{k,n}) + d_H(\Omega_\infty^{k,n-1}).$$

We have the corresponding decomposition

$$\Omega_\infty^{n+k} = \text{Ker } e_k \oplus E_k, \tag{30}$$

where

$$\text{Ker } e_k = d_H(\Omega_\infty^{k,n-1}) \oplus \left(\bigoplus_{m>0} \Omega_\infty^{k+m,n-m} \right). \tag{31}$$

It is readily observed that $\varepsilon_k = \tau_k \circ d$ on $\Omega_\infty^{k-1,n}$. The diagram (29) is derived from the variational bicomplex (see, e.g., [8, 15]). Its first row is a subcomplexes of the infinite-order De Rham complex (11).

Since the diagram (29) coincides with the diagram (22) at exterior forms of degree $\leq n$, the corresponding exact sequence of cohomology groups for the diagram (29) differs from the exact sequence (23) starting from the terms after $H^n(\Omega_\infty^n)$. If $Y \rightarrow X$ is an affine bundle, then $H^{>n}(\Omega_\infty^*) = 0$ and the exact sequence at these terms breaks into the short sequences

$$0 \longrightarrow H^n(X) \xrightarrow{h_0^*} H_{\text{var}}^n \xrightarrow{d^*} H(\text{Ker } e_1) \xrightarrow{\text{in}^*} 0, \tag{32}$$

$$0 \longrightarrow H_{\text{var}}^{n+k} = H(e_{k+1}) \longrightarrow 0, \quad k > 0. \tag{33}$$

Using these exact sequences and the definition (31) of $\text{Ker } e_k$, one can find the cohomology groups $H_{\text{var}}^{\geq n}$ of the variational complex (1). Recall that cocycles in $\text{Ker } e_k$ are closed forms $\phi \in \text{Ker } e_k$, while the coboundaries are exact forms $d\xi$ where $\xi \in \text{Ker } e_{k-1}$.

LEMMA 14. *The cohomology group $H(e_1)$ is trivial.*

Proof. The obstruction for the cohomology group $H(e_1)$ to be trivial are the elements of the overlap $d\Omega_\infty^{0,n} \cap d_H\Omega_\infty^{k,n-1}$, i.e., the forms $\phi \in \Omega_\infty^{0,n}$ such that $d_V\phi = d_H\varepsilon$, i.e., $\phi \in \text{Ker } d_Hd_V$. By virtue of Lemma 12, such a form is given by the sum (28) which reads

$$\phi = d_H\psi + \varphi.$$

Then, $d\phi = -dd_V\psi$ is an exact form. □

It follows that the short exact sequence (32) leads to the isomorphism

$$H^n(X) = H_{\text{var}}^n. \tag{34}$$

In other words, the cohomology group of variationally trivial Lagrangians on an affine bundle modulo d_H -exact forms coincides with the cohomology group $H^n(X)$ of the base manifold X . This states Theorem 3. The isomorphism (34) also leads to the isomorphism

$$H^n(d_H)/H^n(X) = \varepsilon_1(\Omega_\infty^{0,n}),$$

where $H^n(d_H)$ is the n th-cohomology group of the horizontal complex (2). It generalizes the local equality (3).

One can extend Lemma 14 to higher cohomology groups $H(e_k)$ as follows.

PROPOSITION 15. *The obstruction for the cohomology group $H(e_{k+1})$ and, consequently, the cohomology group H_{var}^{n+k} to be trivial is the second cohomology group of the complex (27) at the term $H^k(d_H)$.*

The result follows from the decomposition (30).

References

1. M.Abbati and A.Manià, On differential structure for projective limits of manifolds, *J. Geom. Phys.*, **29**, 35-63 (1999).
2. I.Anderson, Introduction to the variational bicomplex, *Contemp. Math.*, **132**, 51-74 (1992).
3. I.Anderson, *The Variational Bicomplex*, Academic Press, Boston, 1994.
4. M.Bauderon, Differential geometry and Lagrangian formalism in the calculus of variations, in *Differential Geometry, Calculus of Variations, and their Applications*, Lecture Notes in Pure and Applied Mathematics, **100** (Marcel Dekker, Inc., N.Y., 1985), p. 67-82.

5. R.Bott and L.Tu, *Differential Forms in Algebraic Topology*, Springer-Verlag, Berlin, 1982.
6. P.Dedecker and W.Tulczyjew, Spectral sequences and the inverse problem of the calculus of variations, in *Differential Geometric Methods in Mathematical Physics*, Lect. Notes in Mathematics, **836**, Springer-Verlag, Berlin, 1980, pp. 498-503.
7. L.Dickey, On exactness of the variational bicomplex, *Contem. Math.*, **132**, 307-315 (1992).
8. G.Giachetta, L.Mangiarotti and G.Sardanashvily, *New Lagrangian and Hamiltonian Methods in Field Theory*, World Scientific, Singapore, 1997.
9. I.Krasil'shchik, V.Lychagin and A.Vinogradov, *Geometry of Jet Spaces and Non-linear Partial Differential Equations*, Gordon and Breach, Glasgow, 1985.
10. D.Krupka, J.Musilova, Trivial Lagrangians in field theory, *Diff. Geom. Appl.*, **9**, 293-305 (1998); Erratum, **10**, 303 (1999).
11. S.Mac Lane, *Homology* (Springer-Verlag, Berlin, 1967).
12. W.Massey, *Homology and Cohomology Theory*, Marcel Dekker, Inc., N.Y., 1978.
13. F.Takens, Symmetries, conservation laws and variational principles, in *Geometry and Topology*, Lect. Notes in Mathematics, **597**, Springer-Verlag, Berlin, 1977, pp. 581-604.
14. F.Takens, A global version of the inverse problem of the calculus of variations, *J. Diff. Geom.*, **14**, 543-562 (1979).
15. W.Tulczyjew, The Euler-Lagrange resolution, in *Differential Geometric Methods in Mathematical Physics*, Lect. Notes in Mathematics, **836** (Springer-Verlag, Berlin, 1980), pp. 22-48.